

List of publications

Original articles

1. Schmerzhaftes Myoarthropathien des Kausystems-evidenzbasierte Diagnostik. Türp JC, Schindler HJ, **Bartzela T**. *Kieferorthop* 2005;19:173-81.
2. Long-term stability of unilateral posterior crossbite correction. **Bartzela T**, Jonas I, E. *Angle Orthod* 2007;77:237-43.
3. Load-deflection characteristics of the superelastic NiTi orthodontic wires. **Bartzela T**, Senn C, Wichelhaus A. *Angle Orthod* 2007;77:991-7.
4. Tooth agenesis patterns in bilateral cleft lip and palate patients. **Bartzela T**, Carels C, Bronkhors E, Rønning E, Rizell S, Kuijpers-Jagtman AM. *Eur J Oral Sci* 2009;118:47-52.
5. [Associated congenital anomalies in patients with bilateral cleft lip and palate]. Geassocieerde aangeboren aandoeningen bij bilaterale cheilo-gnato-palatoschisis. Kouwenberg M., Draaisma JM., Kuijpers-Jagtman AM., **Bartzela T**. *Nederlands Tijdschrift Kindergeneesk* 2010;78:15-20.
6. A longitudinal three-center study of dental arch relationship in patients with bilateral cleft lip and palate. **Bartzela T**, Katsaros C, Shaw W, Rønning E, Rizell S, Bronkhorst E, Okada- Ozawa T, Pinheiro F, Dominguez-Gonzales S, Hagberg C, Semb G, Kuijpers-Jagtman AM. *Cleft Palate Craniofac J* 2010;47:167-74.
([2011 Samuel Berkowitz: Long-term outcomes study award](#))
<http://www.cleftline.org/healthcare-professionals/awards-scholarships/#berkowitz>
7. A two-centre study on facial morphology in patients with complete bilateral cleft lip and palate at nine years of age. **Bartzela T**, Katsaros C, Bronkhorst E, Rizell S, Halazonetis D, Kuijpers-Jagtman AM. *Int J Oral Maxillofac Surg*. 2011;40:782-9.
8. Comparison of two scoring systems for evaluation of dental arch relationships in patients with complete bilateral cleft lip and palate. **Bartzela T**, Leenarts C, Bronkhorst E, Borstlap W, Katsaros C, Kuijpers-Jagtman AM. *Cleft Palate Craniofac J* 2011;48:455-61.
9. Dental agenesis patterns of permanent teeth in Crouzon syndrome patients. Stavropoulos D, **Bartzela T**, Mohlin B, Hagberg C. *Eur J Oral Sci* 2011;119:198-203.

10. Tooth agenesis patterns in bilateral cleft lip/palate (BCLP) indicate possible common genetic pathways. Martin MD. *J Evid Based Dent Pract* 2010;10:252-3.
11. Photographs of dental casts or digital models: rating dental arch relationships in bilateral cleft lip and palate. Leenarts C, **Bartzela T**, Bronkhorst E, Semb G, Shaw W, Katsaros C, Kuijpers-Jagtman AM. *Int J Oral Maxillofac Surg.* 2012;41:180-5.
12. Stavropoulos D, **Bartzela T**, Mohlin B, Hagberg C. Dental agenesis patterns of permanent teeth in Apert syndrome patients. *Swed Dent J* 2011;35:195-211.
13. A longitudinal three-center study of craniofacial morphology at 6 and 12 years of age in patients with complete bilateral cleft lip and palate. **Bartzela T**, Katsaros C, Rønning E, Rizell S, Semb G, Bronkhorst E, Halazonetis D, Kuijpers-Jagtman AM. *Clin Oral Invest.* 2012; 16:1313-24. 14.
14. Tooth agenesis patterns in unilateral cleft lip and palate patients. Bartzela T, Carels C, Bronkhors E, Kuijpers-Jagtman AM. *Arch Oral Biol* 2013; 58:596-602.

Review Articles

15. Medication effects on the rate of orthodontic tooth movement: a systematic literature review. **Bartzela T**, Türp JC, Motschall E, Maltha JC. *Am J Orthod Dentofacial Orthop* 2009;135:16-26. Written examination of American Board of Orthodontics since 2010
<http://www.americanboardortho.com/orthodontic-professionals/about-board-certification/written-examination/journal-articles/>
16. [Eruption guidance appliances in orthodontics]. Eruptiegeleidende apparatuur in de orthodontie. Reukers HAJ, **Bartzela T**. *Ned Tijdsch Tandheelkd* 2008;115:125- 31.